



**International Journal of Computational Economics and Econometrics**

ISSN online: 1757-1189 - ISSN print: 1757-1170

<https://www.inderscience.com/ijcee>

---

**Identifying trend nature in time series using autocorrelation functions and stationarity tests**

M. Boutahar, M. Royer-Carenzi

**DOI:** [10.1504/IJCEE.2023.10056902](https://doi.org/10.1504/IJCEE.2023.10056902)

**Article History:**

Received:	05 August 2021
Last revised:	06 December 2022
Accepted:	25 March 2023
Published online:	20 December 2023

---

## Identifying trend nature in time series using autocorrelation functions and stationarity tests

---

M. Boutahar\* and M. Royer-Carenzi

CNRS,  
Centrale Marseille,  
Aix Marseille University,  
I2M, UMR 7373,  
Marseille, France  
Email: mohamed.boutahar@univ-amu.fr  
Email: manuela.royer-carenzi@univ-amu.fr  
\*Corresponding author

**Abstract:** Time series non-stationarity can be detected thanks to autocorrelation functions. But trend nature, either deterministic or either stochastic, is not identifiable. Strategies based on Dickey-Fuller unit root-test are appropriate to choose between a linear deterministic trend or a stochastic trend. But all the observed deterministic trends are not linear, and such strategies fail in detecting a quadratic deterministic trend. Being a confounding factor, a quadratic deterministic trend makes a unit root spuriously appear. We provide a new procedure, based on Ouliaris-Park-Phillips unit root test, convenient for time series containing polynomial trends with a degree higher than one. Our approach is assessed based on simulated data. The strategy is finally applied on two real datasets: money stock in USA and on CO<sub>2</sub> atmospheric concentration. Compared with Dickey-Fuller diagnosis, our strategy provides the model with the best performances.

**Keywords:** time series; stationarity; autocorrelation functions; unit root tests; Dickey-Fuller; KPSS; OPP test; trend detection; deterministic or stochastic trend; spurious unit root.

**Reference** to this paper should be made as follows: Boutahar, M. and Royer-Carenzi, M. (2024) 'Identifying trend nature in time series using autocorrelation functions and stationarity tests', *Int. J. Computational Economics and Econometrics*, Vol. 14, No. 1, pp.1–22.

**Biographical notes:** M. Boutahar is an Associate Professor in Aix-Marseille University since 1992, and a researcher in Applied Mathematics in the ALEA research team of the Institute of Mathematics of Marseille, UMR 7373 since 2006. His research interests: time series analysis; statistical tests of breaks (mean, variance, distribution, tail index); long memory models, nonlinear and non-stationary models (estimates, tests, forecasts); multivariate extreme value theory.

M. Royer-Carenzi is an Associate Profesor since 2004, and a Fellow of I2M Laboratory in the Aix-Marseille University since 2007. She started working on backward stochastic differential equations, with a PhD in the Rennes

University, France, in 2003. Thanks to collaborations with biologists and actuaries, she gets interested in applied statistics, with main applications in ecology, in epidemiology and in phylogeny.

---

## 1 Introduction

Time series non-stationarity can originate from various sources: either from a trend component or from a seasonal or even a cyclical component. In this paper, we will be interested in the non-stationarity caused by a trend. There are two kinds of trends: either a deterministic trend which can be modelled by some function of time (polynomial trend is generally considered), or a stochastic trend which presents unit roots. Deterministic and stochastic trends are two specific models suggested by Nelson and Plosser (1982).

$$\text{Deterministic trend (\textbf{Det}, \mathbf{d})} \quad Z_t = a_0 + a_1 t + \dots + a_d t^d + B_t \quad (1)$$

$$\text{Stochastic trend (\textbf{Sto}, \mathbf{d})} \quad \Delta^d(Z_t) = B_t, \quad (2)$$

where  $d$  is an integer and  $a_d \neq 0$ . Moreover,  $L$  is the backshift operator, i.e.,  $LZ_t = Z_{t-1}$ , so that the 1-lag difference operator  $\Delta$  can be expressed as

$$\Delta(Z_t) = (1 - L)(Z_t) = Z_t - Z_{t-1}.$$

Finally  $(B_t)_t$  is a moving average process

$$B_t = \sum_{j \in \mathbb{Z}} b_j \mathcal{E}_{t-j}, \quad (\text{H1})$$

where  $(\mathcal{E}_t)_t$  is a sequence of identically distributed and independent centred variables, such that

$$E(\mathcal{E}_t^{2k}) < \infty, \text{ for some } k \geq 2, \quad (\text{H2})$$

and where parameters  $b_j$  ( $j \in \mathbb{Z}$ ) satisfy

$$\begin{aligned} \sum_{j \in \mathbb{Z}} |b_j| < \infty & \quad \sum_{j \in \mathbb{Z}} b_j \neq 0 & \quad \sum_{j \in \mathbb{Z}} b_j^2 |j| < \infty. \\ (\text{H3-a}) & \quad (\text{H3-b}) & \quad (\text{H3-c}) \end{aligned} \quad (\text{H3})$$

We denote  $(B_t)_t$  as **(SN)**, for *stationary noise*. It is well known that causal and invertible ARMA processes satisfy Hypotheses (H1) and (H3). When  $B_t = \mathcal{E}_t$ , it is called **(WN)** for *white noise*. In this case, the associated models defined in equations (1) and (2), are referred as **(Det<sub>W</sub>, d)** and **(Sto<sub>W</sub>, d)**.

When modelling time series, specially for macroeconomic and financial data, it is very important to identify the nature of the trend: either deterministic or stochastic. Indeed, every type of trend induces specific behaviours, that we can illustrate with moments properties. Let us consider a **(Det, 1)** then  $\mathbb{E}(Z_t) = a_0 + a_1 t$ , and  $\text{var}(Z_t) = \sigma_B^2$ , providing a stationary variance but a non-stationary mean. On the contrary, under

a **(Sto, 1)** model with  $Z_0 = 0$ , we have  $\mathbb{E}(Z_t) = 0$ , and  $\text{var}(Z_t) = t\sigma_B^2$ , providing the opposite feature. Thus the source of non-stationarity differs with trend nature. Consequently identifying the correct trend is fundamental. Globally, time series with a deterministic trend always revert to the trend in the long run (the effects of shocks are eventually eliminated); and the prediction intervals have constant width. On the contrary, time series with a stochastic trend never recover from shocks to the mean (the effects of shocks are permanent); and the forecast confident intervals grow with the horizon. Several authors (see for instance Chan et al., 1977; Nelson and Kang, 1981) studied the consequences of an inappropriate modelling choice, underlying the importance of developing procedures able to produce a reliable classification.

Note that non-stationarity coming from a periodical trend (seasonal or even a cyclical) can also be considered. One can consider the following models:

$$\text{Deterministic seasonality } Z_t = \mathbf{S}_t + Y_t, \quad (3)$$

$$\text{Stochastic seasonality } \Delta_r^D(Z_t) = Y_t, \quad (4)$$

where  $Y_t$  has a trend, which can be modelled by either model (1) or (2). Moreover, in (3), we define  $\mathbf{S}_t = S_i$  if  $(t \bmod r) = i$ , and  $S_1, \dots, S_r$  are the seasonal components. And in (4), we consider the seasonal difference operator ( $r$ -lag difference operator)

$$\Delta_r(Z_t) = (1 - L^r)(Z_t) = Z_t - Z_{t-r},$$

where  $r$  is the season assumed to be known and  $D$  is an integer.

Autocorrelation function properties have been widely studied, and are helpful to specify accurate models for stationary time series. The theoretical autocorrelation function at lag  $h$  ( $|h| < n$ ) is estimated from data  $(Z_1, \dots, Z_n)$  with the random variable

$$\Xi(h) = \frac{\sum_{t=1}^{n-h} (Z_{t+h} - \bar{Z})(Z_t - \bar{Z})}{\sum_{t=1}^n (Z_t - \bar{Z})^2}, \quad (5)$$

where

$$\bar{Z} = \frac{\sum_{t=1}^n Z_t}{n},$$

is the random mean. We show that autocorrelation functions also have interesting properties in our framework, since they permit to identify the presence of a trend in a time series. Nevertheless it cannot be used to distinguish between either a deterministic or a stochastic trend.

Dickey and Fuller (1979) developed a unit root test that is an essential tool in time series modelling. The authors considered the following models:

$$\mathbf{M}_1 : Z_t = \rho Z_{t-1} + \mathcal{E}_t$$

$$\mathbf{M}_2 : Z_t = a_0 + \rho Z_{t-1} + \mathcal{E}_t$$

$$\mathbf{M}_3 : Z_t = a_0 + a_1 t + \rho Z_{t-1} + \mathcal{E}_t,$$

where  $(\mathcal{E}_t)_t$  is a white noise. Dickey-Fuller procedure permits to test the null hypothesis of a unit root ( $\rho = 1$ ) against the alternative hypothesis of a stationary AR(1) model without drift (resp. with drift, resp. with trend), accordingly to model  $\mathbf{M}_1$  (resp.  $\mathbf{M}_2$ , resp.  $\mathbf{M}_3$ ). Note that a time series with a (linear) deterministic trend, as defined in  $(\mathbf{Det}_W, 1)$ , is included in  $\mathbf{M}_3$  model, by taking  $\rho = 0$ . Consequently, under a (linear) deterministic trend, Dickey-Fuller unit-root test, led in  $\mathbf{M}_3$  framework, usually rejects the null hypothesis. On the other hand,  $(\mathbf{Sto}_W, 1)$ -time series are included in  $\mathbf{M}_1$  model, with  $\rho = 1$ , so that  $(H_0)$ , tested in  $\mathbf{M}_1$  framework, is usually not rejected, accordingly with the rejection rate.

Several strategies based on Dickey-Fuller unit root-test have been developed (Dickey et al., 1986; Perron, 1988; Dolado et al., 1990), and they mostly appear to be appropriate to choose between a linear deterministic trend or a stochastic trend. But all the observed deterministic trends are not linear, trends with higher degrees can be involved. In this case, Ertur (1998) noticed that the usual strategies fail in detecting a quadratic deterministic trend. Indeed, under a quadratic trend for instance, Dickey-Fuller test generally concludes to the presence of a (spurious) unit root, even using model  $\mathbf{M}_3$  that allows a linear trend.

In this paper, we aim to include polynomial trends with a degree higher than one. Let us introduce the general model:

$$\mathbf{M}_{3,d} : Z_t = a_0 + a_1 t + \dots + a_d t^d + \rho Z_{t-1} + B_t \quad (a_d \neq 0).$$

In Ouliaris et al. (1989), the authors developed a test that corrects the bias caused by high degree trend when testing for a unit root. We included this test in a strategy, that correctly identifies either a deterministic trend or a stochastic one.

In Section 2, we detail autocorrelation functions convergence when time series are driven by a trend, either deterministic or stochastic. This result permits to detect the presence of a trend, but without precise identification. We also study existing strategies based on Dickey-Fuller tests, and analyse their performance in classifying models  $(\mathbf{Det}_W, 1)$ ,  $(\mathbf{Det}_W, 2)$ ,  $(\mathbf{Sto}_W, 1)$  and  $(\mathbf{Sto}_W, 2)$ . Note that we also consider the simplest model  $(\mathbf{WN})$  as a null-model. Next, we provide a new strategy, based on trend diagnosis tests (TDT), theoretically able to identify trends type, even when higher-degrees  $d$  are considered. In this paper we study the asymptotic behaviour of autocorrelation functions only for models (1) and (2). Similar results can be obtained for models (3) and (4) but are out of the scope of this paper. However, to deal with seasonal data (monthly, quarterly or biannual) one can extract the seasonal component first and then apply our proposed strategy. In Section 3, we ran simulations of processes generated by models  $(\mathbf{WN})$ ,  $(\mathbf{Det}_W, 1)$ ,  $(\mathbf{Det}_W, 2)$ ,  $(\mathbf{Sto}_W, 1)$  such as  $(\mathbf{Sto}_W, 2)$  and we applied the various tests of stationarity studied previously. It appears that the Dickey-Fuller tests fail in the presence of a degree 2 trend. On the other hand, the tests included in our TDT strategy behave as expected on simulations even when  $(\mathbf{WN})$  is replaced by a  $(\mathbf{SN})$  underlying process. But because of type I of type II errors associated with the different tests, the simulations also reveal responses that are complementary to the theoretical one. This provides a modelling suggestion when the series does not produce one of the theoretical responses. Finally, in Section 4, we apply our strategy on real data such as money stock in the USA and also on the  $\text{CO}_2$  atmospheric concentration, that includes a monthly seasonal component. All the functions are implemented in **R** language, and they are available at the website:

- <http://www.i2m.univ-amu.fr/perso/manuela.royer-carezzi/AnnexesR.TrendTS/TrendTS.html>

## 2 Trend detection and nature identification

### 2.1 Detection with autocorrelation functions

It is essential to start by plotting the graph of the series in order to visualise its evolution, but the presence or absence of a trend is sometimes difficult to detect from the plot. Autocorrelation appears to be a powerful tool in the detection of a trend insofar as its behaviour is specific in the presence of a trend. We recall that theoretical autocorrelations  $\rho(h) = \text{cor}(Z_t, Z_{t+h})$  are well-defined only if process  $(Z_t)_t$  is square-integrable and stationary. However the associated random variables  $\Xi(h)$ , as defined in equation (5), can be computed for any observed time series  $(Z_1, \dots, Z_n)$ .


#### 2.1.1 Autocorrelation functions behaviour for (WN)-time series

*Theorem 2.1:* Let  $(Z_t)_t$  be a white noise. Then

$$\sqrt{n} {}^t(\Xi(1), \Xi(2), \dots, \Xi(r)) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \mathcal{N}_r({}^t(0, \dots, 0), Id_r),$$

where  ${}^tv$  denotes the transpose of vector  $v$ , and  $Id_r$  is the identity  $r \times r$ -matrix.

Theorem 2.1 is a particular case of Theorems 7.2.1. or 7.2.2. in Brockwell and Davis (1991), that require Hypotheses (H1), (H3-a) and either Hypothesis (H2) or (H3-c). Thus, we get that the random variables  $\Xi(1), \Xi(2), \dots, \Xi(r)$  are asymptotically independent and identically distributed as Gaussian random variables with zero mean and variance  $1/n$ . Consequently, for any fixed lag  $h = 1, \dots, r$ , for large  $n$ , the sample autocorrelation function  $\sqrt{n} \hat{\rho}(h)$  is expected to be a realisation of a standard Gaussian, that is to be valued in the interval  $[-1.96, 1.96]$ , with 95% coverage. Thus, sample autocorrelation functions (acf) are used to assess for white noise. But even when the underlying process is a white noise, several autocorrelations among  $\hat{\rho}(1), \dots, \hat{\rho}(r)$  may lie out of the interval  $[-1.96/\sqrt{n}, 1.96/\sqrt{n}]$ . The asymptotic independence property for variables  $\Xi(h)$  implies that, when sample size  $n$  is large, the number of observed autocorrelation functions out of this interval behaves as a binomial  $\mathcal{B}(r, 0.05)$  distribution. We take into account the multiple testing paradigm, on the one hand by incorporating the binomial exact test and on the other hand by incorporating a global testing procedure, using Sidak correction. As an example, we simulate a white noise, and compute sample autocorrelation functions. Figure 1 shows that three values lie out of the interval  $[-1.96/\sqrt{n}, 1.96/\sqrt{n}]$ , plotted with blue dashed lines. But binomial exact test (p-value = 0.1159) confirms that such a number remains consistent with white noise hypothesis. Moreover, a second set of interval, computed with Sidak correction, is provided, and plotted with red dotted lines. If at least one sample autocorrelation function lies out this global interval, then white noise hypothesis is rejected. In our simulation, white noise diagnosis is confirmed, both by graphics and by binomial's test.


**Figure 1** Sample autocorrelation functions for a white noise simulation (see online version for colours)

### 2.1.2 Autocorrelation functions for **(Det, d)**-time series

**Theorem 2.2:** Let  $(Z_t)_t$  be a stochastic process such that  $Z_t = \sum_{j=0}^d a_j t^j + B_t$ , where  $a_d \neq 0$  and  $(B_t)_t$  is **(SN)** satisfying Hypotheses (H1) to (H3). Then

$$\Xi(h) \xrightarrow[n \rightarrow +\infty]{\mathbf{P}} 1, \quad \forall h \neq 0.$$

Proof is given in Appendix A. Figure 2(a) illustrates the slow decreasing behaviour of sample autocorrelation functions when time series are driven by a deterministic **(Det, 2)** trend.

**Figure 2** Sample autocorrelation functions for simulations with either (a) a deterministic or (b) a stochastic trend (see online version for colours)

### 2.1.3 Autocorrelation functions for $(\mathbf{Sto}_W, \mathbf{d})$ -time series

**Theorem 2.3:** Let  $(Z_t)_t$  be a stochastic process such that  $\Delta^d(Z_t) = \mathcal{E}_t$ , with  $Z_t = 0$ , for any  $t \leq 0$  and  $(B_t)_t$  is **(SN)** satisfying Hypotheses (H1) to (H3). Then

$$\Xi(h) \xrightarrow[n \rightarrow +\infty]{\mathbf{P}} 1, \quad \forall h \neq 0.$$

Proof is given in Appendix B. Figure 2(b) illustrates the slow decreasing behaviour of sample autocorrelation functions when time series are driven by a stochastic  $(\mathbf{Sto}_W, \mathbf{2})$  trend.

## 2.2 Trend-nature identification with stationarity tests

Previously, we showed that the autocorrelation functions, computed from variables  $Z_1, \dots, Z_n$ , have a particular asymptotic behaviour in presence of a trend. But Figure 2 illustrates that the behaviour is similar either for a deterministic or a stochastic trend. Consequently, a deeper study has to be led in order to specify the type of trend.

### 2.2.1 Data generating process

We explore stationarity tests diagnosis according to several data generating processes (DGPs), and we will also explore the associated differenced processes. We set  $(B_t)_t$  a stationarity noise constructed from a white noise  $(\mathcal{E}_t)_t$  with variance  $\sigma_{\mathcal{E}}^2$ . We consider processes  $(Z_t)_t$  successively driven by one of the following models:

$$(\mathbf{SN}) \quad Z_t = B_t \implies \Delta(Z_t) = B_t - B_{t-1}, \quad (6)$$

$$(\mathbf{Det}, \mathbf{1}) \quad Z_t = a_0 + a_1 t + B_t \implies \Delta(Z_t) = a_1 + B_t - B_{t-1}, \quad (7)$$

$$(\mathbf{Det}, \mathbf{2}) \quad Z_t = a_0 + a_1 t + a_2 t^2 + B_t \implies \Delta(Z_t) = a_1 + 2a_2 t + B_t - B_{t-1}, \quad (8)$$

$$(\mathbf{Sto}, \mathbf{1}) \quad Z_t = Z_{t-1} + B_t \implies \Delta(Z_t) = B_t, \quad (9)$$

$$(\mathbf{Sto}, \mathbf{2}) \quad Z_t = 2Z_{t-1} - Z_{t-2} + B_t \implies \Delta(Z_t) = \Delta(Z_{t-1}) + B_t. \quad (10)$$

Some of the previous processes are stationary while others are not. We recall that a process containing a unit root is not stationary. Thus models **(Sto, 1)** and **(Sto, 2)** do contain a unit root and hence are not stationary, whereas models **(Det, 1)** and **(Det, 2)** are not stationary although they do not contain any unit root. Table 1 indicates which of these series or differentiated series is stationary or has a unit root.

### 2.2.2 Dickey-Fuller tests

In Dickey and Fuller (1979), the authors introduced unit root tests adapted to models  $\mathbf{M}_1, \mathbf{M}_2$  and  $\mathbf{M}_3$ . Every time the null and the alternative hypothesis are mathematically expressed in the same way:

$$(\underline{H}_0) : \rho = 1 \text{ against } (\underline{H}_1) : |\rho| < 1.$$



**Table 1** Stationarity and presence of a unit root for processes generated by models **(SN)**, **(Det, 1)**, **(Det, 2)**, **(Sto, 1)** such as **(Sto, 2)**, and for the associated differenced processes

Property	DGP <sup>a</sup>				
	(SN)	(Det, 1)	(Det, 2)	(Sto, 1)	(Sto, 2)
Stationarity for $Z_t$	Yes	No	No	No	No
Stationarity for $\Delta(Z_t)$	Yes	Yes	No	Yes	No
Unit root for $Z_t$	No	No	No	Yes	Yes
Unit root for $\Delta(Z_t)$	No	No	No	No	Yes

Note: <sup>a</sup>DGP.

But the alternative hypothesis interpretation depends on the considered model.

$(\mathbf{M}_1)(\underline{H}_1)$ :  $(Z_t)_t$  is a stationary and centred AR(1) process;

$(\mathbf{M}_2)(\underline{H}_1)$ :  $(Z_t)_t$  is a stationary not-centred AR(1) process;

$(\mathbf{M}_3)(\underline{H}_1)$ :  $(Z_t)_t$  is (linear-)trend stationary (TS).

And test statistics do not have the same expression, possibly leading to opposing conclusions, even on the same data. To distinguish between the null or the alternative hypothesis, one has to use the suitable test statistics, adapted to every model.

In Dickey and Fuller (1979), the authors also developed joined tests:

Test  $\Phi_1$  under  $(\mathbf{M}_2)(\underline{H}_0)$ :  $(a_0, \rho) = (0, 1)$ ;

Test  $\Phi_2$  under  $(\mathbf{M}_3)(\underline{H}_0)$ :  $(a_0, a_1, \rho) = (0, 0, 1)$ ;

Test  $\Phi_3$  under  $(\mathbf{M}_3)(\underline{H}_0)$ :  $(a_0, a_1, \rho) = (a_0, 0, 1)$ .

Models **(Sto<sub>W</sub>, 1)** and **(Sto<sub>W</sub>, 2)** do contain a unit root and hence are not stationary, whereas models **(Det<sub>W</sub>, 1)** and **(Det<sub>W</sub>, 2)** are not stationary although they do not contain any unit root. Logically, Dickey-Fuller test should not reject the null hypothesis for almost realisations driven from models **(Sto<sub>W</sub>, 1)** and **(Sto<sub>W</sub>, 2)**, precisely with a rate  $(1 - \alpha)\%$ , where  $\alpha$  stands for the significance level. Reciprocally, under **(Det<sub>W</sub>, 1)** and **(Det<sub>W</sub>, 2)** models, the null hypothesis should be rejected.

### 2.2.3 OPP stationarity test

In Ouliaris et al. (1989), the authors generalised Dickey-Fuller unit-root tests  $\rho$  to models  $\mathbf{M}_{3,d}$  with polynomial trends, where  $d = 2, 3, 4$  or  $5$ . We denote this general test by OPP. Note that the invariance principle for partial sums, required in OPP test, applies to **(SN)** stationary linear processes satisfying Hypotheses (H1) to (H3).

From Table 1, we guess that OPP test should reject the null hypothesis only for models **(Det, 1)** and **(Det, 2)**, vice versa for **(Sto, 1)** and **(Sto, 2)**. Applied to the differentiated series, OPP test should not reject the null hypothesis only for realisations initially driven from models **(Sto, 2)**.

### 2.2.4 KPSS stationarity test

In Kwiatkowski et al. (1992), the authors developed another type of stationarity test, associated to an underlying **(SN)**. Contrary to Dickey-Fuller and OPP tests, KPSS

test takes the presence of unit root as the alternative hypothesis, and the stationarity as the null hypothesis. Actually, KPSS test can consider as null-hypothesis either level-stationarity or trend-stationarity (stationarity around a linear deterministic trend). However, there is no extension in the literature of KPSS test considering stationarity around a polynomial deterministic trend.

Note that none model is stationary, except the simplest model (SN). Hence KPSS test does not appear to be a good candidate for trend-nature identification. Nonetheless, KPSS test may reveal heterogeneous behaviours when applied to the differentiated series  $\Delta(Z_t)$ . Indeed, processes  $(Z_t)$  driven from models (Det, 1) and (Sto, 1) do become stationary as soon as they are differentiated.

Here we will use the level stationarity. Note that KPSS test does not include the case of series with non-invertible noise, as is the case for the (SN) and (Det, 1) differentiated series. But it is possible to determine the nature of the convergence of the test statistic in these particular cases (see Appendix C.1), and we even identify the behaviour of the test statistic under an alternative such as a (Det, 1) or a (Det, 2) process (see Appendix C.2).

## 2.2.5 TDT strategy

### 2.2.5.1 Description and theoretical diagnosis

We introduce a strategy to identify trend nature by applying the following tests successively:

- 1 OPP test to series  $Z_t$
- 2 OPP test to series  $\Delta(Z_t)$
- 3 KPSS test to series  $Z_t$
- 4 KPSS test to serie  $\Delta(Z_t)$ .

We call TDT the set of responses to tests 1 to 4 computed on a time series. Let us denote by **Null**, the case where the null hypothesis cannot be rejected, and by **Alt** otherwise. From equations (6) to (10), we provide the theoretical response obtained by applying the TDT strategy for the following five models:

$$(\text{SN}) \text{ theoretical response: } \text{Alt/Alt/Null/Null}, \quad (11)$$

$$(\text{Det}, 1) \text{ theoretical response: } \text{Alt/Alt/Alt/Null}, \quad (12)$$

$$(\text{Det}, 2) \text{ theoretical response: } \text{Alt/Alt/Alt/Alt}, \quad (13)$$

$$(\text{Sto}, 1) \text{ theoretical response: } \text{Null/Alt/Alt/Null}, \quad (14)$$

$$(\text{Sto}, 2) \text{ theoretical response: } \text{Null/Null/Alt/Alt}. \quad (15)$$

For example the (SN) model does not contain unit root, and is stationary (see Table 1). So is  $\Delta(\text{SN})$ . Hence the OPP test will accept the alternative of no unit root for (SN) and  $\Delta(\text{SN})$ , then we have the two responses **Alt/Alt/**. On the other hand the KPSS test will accept the null hypothesis of stationarity for (SN) and  $\Delta(\text{SN})$ ; then we have the two responses **/Null/Null**. Therefore the theoretical response of the TDT strategy for the (SN) model will be **Alt/Alt/Null/Null**. A similar explanations can be made for the other models to obtain the theoretical responses of the TDT strategy.

### 2.2.5.2 More diagnosis expected for simulations

Note that, when applying the TDT strategy to a given DGP, there are 16 possible responses; one response is the theoretical one, as given in equations (11)–(15), whereas the other responses are not the accurate ones. For example if the DGP is (SN) model, then the expected response will be the theoretical one **Alt/Alt/Null/Null**, and the wrong responses are **w/x/y/z** where **w, x, y, z** can take either **Alt** or **Null**, except the case **w = Alt, x = Alt, y = Null, z = Null**. We insist on the fact that when applying the TDT procedure on a simulation, the observed diagnosis will not systematically be the theoretical one. Indeed, a statistical test, even a well-calibrated and powerful one, produces erroneous responses, in proportion to type I and type II errors. Most often, the power of a test is difficult to calculate, but when the null hypothesis is verified, the null hypothesis is still rejected in  $\alpha\%$  of cases, where  $\alpha$  is the nominal level.

- Thus when we consider for example a (WN) model, only the last two tests 3 and 4 are performed under the null hypothesis of stationarity. But we point out that KPSS test was developed under a white noise hypothesis, that is clearly convenient for the initial series, but no longer valid for the differentiated series. Consequently, KPSS behaviour in test 3 will be convenient, whereas KPSS behaviour in test 4, i.e., on the differentiated series, which is nothing else than a non-invertible MA(1) process, will not necessarily be in adequacy with the proportions  $\alpha$  and  $1 - \alpha$ . Actually, we proved in Appendix C.1, equation (C2), that KPSS test will never reject the null hypothesis, as  $n$  tends to  $\infty$ , for series  $(\Delta(Z_t))_t$  when  $(Z_t)_t$  is (WN). Consequently, we expect, except for size distortion effect, that

$$\begin{aligned}
 95\% \times 100\% &= 95\% && \text{(WN)-simulations with response -/-/Null/Null} \\
 95\% \times 0\% &= 0\% && \text{(WN)-simulations with response -/-/Null/Alt} \\
 5\% \times 100\% &= 5\% && \text{(WN)-simulations with response -/-/Alt/Null} \\
 5\% \times 0\% &= 0\% && \text{(WN)-simulations with response -/-/Alt/Alt}
 \end{aligned} \tag{16}$$

- In the same way, for a (**Det<sub>W</sub>, 1**), none of the tests 1 to 3 is performed under the null hypothesis. And for test 4, it is the same as applying the KPSS test on a non-invertible MA(1) process. We recall that we proved in Appendix C.1, equation (C2), that KPSS test will never reject the null hypothesis, as  $n$  tends to  $\infty$ , for series  $(\Delta(Z_t))_t$  when  $(Z_t)_t$  is (**Det<sub>W</sub>, 1**). Moreover, we proved in Appendix C.2, equation (C3), that KPSS test will always reject the null hypothesis, as  $n$  tends to  $\infty$ , for series under a (**Det<sub>W</sub>, 1**) model, meaning that KPSS test has a very high power (probability of rejecting the null when the alternative is true) in this case. Hence both tests 3 and 4 have a tractable behaviour. Consequently, we expect, except for size distortion effect, that

$$\begin{aligned}
 0\% \times 100\% &= 0\% && \text{(Det}_W, 1\text{)-simulations with response -/-/Null/Null} \\
 0\% \times 0\% &= 0\% && \text{(Det}_W, 1\text{)-simulations with response -/-/Null/Alt} \\
 100\% \times 100\% &= 100\% && \text{(Det}_W, 1\text{)-simulations with response -/-/Alt/Null} \\
 100\% \times 0\% &= 0\% && \text{(Det}_W, 1\text{)-simulations with response -/-/Alt/Alt}
 \end{aligned} \tag{17}$$

- For model **(Det<sub>W</sub>, 2)**, none of the tests 1 to 4 is performed under the null hypothesis. But once again, KPSS behaviour is tractable both on  $(Z_t)_t$  and  $(\Delta(Z_t))_t$  processes. Indeed, in Appendix C.2, equation (C3), we showed that KPSS test will always reject the null hypothesis, as  $n$  tends to  $\infty$ , for series  $(Z_t)_t$  and  $(\Delta(Z_t))_t$  when  $(Z_t)_t$  is generated under a **(Det<sub>W</sub>, 2)** model, meaning that KPSS test has a very high power in both cases. Consequently, we expect, except for size distortion effect, that

$$\begin{aligned}
 0\% \times 0\% &= 0\% && \textbf{(Det}_W, \mathbf{2})\text{-simulations with response -/-Null/Null} \\
 0\% \times 100\% &= 0\% && \textbf{(Det}_W, \mathbf{2})\text{-simulations with response -/-Null/Alt} \\
 100\% \times 0\% &= 0\% && \textbf{(Det}_W, \mathbf{2})\text{-simulations with response -/-Alt/Null} \\
 100\% \times 100\% &= 100\% && \textbf{(Det}_W, \mathbf{2})\text{-simulations with response -/-Alt/Alt}
 \end{aligned} \tag{18}$$

- Let us now consider a **(Sto<sub>W</sub>, 1)** process, among tests 1 to 4, only the first and the last one are led under the null hypothesis. Consequently, when  $\alpha = 5\%$ , each of tests 1 and 4 may approximately produce the **Null** response in 95% simulations, and the **Alt** response in the other 5%. This yields to the following theoretical percentages

$$\begin{aligned}
 95\% \times 95\% &= 90.25\% && \textbf{(Sto}_W, \mathbf{1})\text{-simulations with response Null/-/-Null} \\
 95\% \times 5\% &= 4.75\% && \textbf{(Sto}_W, \mathbf{1})\text{-simulations with response Null/-/-Alt} \\
 5\% \times 95\% &= 4.75\% && \textbf{(Sto}_W, \mathbf{1})\text{-simulations with response Alt/-/-Null} \\
 5\% \times 5\% &= 0.25\% && \textbf{(Sto}_W, \mathbf{1})\text{-simulations with response Alt/-/-Alt}
 \end{aligned} \tag{19}$$

- Finally for **(Sto<sub>W</sub>, 2)** process, among tests 1 to 4, only the first and the second one are led under the null hypothesis of the presence of an unit root. Let us point out that OPP test is very efficient to detect single unit roots, but it has not been developed to deal with the case of multiple unit roots. Consequently, OPP behaviour in test 2, i.e., on the differentiated series, which is nothing else than a **(Sto<sub>W</sub>, 1)** process, will be adequate and can be anticipated with proportion  $\alpha$  for **Alt** response, and respectively  $1 - \alpha$  for **Null**; while the answers given on the initial series will not necessarily be in adequacy with these proportions.

### 2.2.5.3 Higher degree trends

Actually, it is possible to detect higher-degree trends, either deterministic **(Det, d)** or stochastic **(Sto, d)**, with  $d = 3, 4, 5$ , by iterating OPP and KPSS tests on the successive differentiated series. More precisely,

- *Step 0:*  
Compute sample autocorrelation functions in order to distinguish between a white noise and a time series with a trend. If the series is driven by a trend, then run the following steps.
- *Step 1:*  
Run OPP test on the given time series.

If the null is rejected, we identify a **(Det, d)** model, otherwise **(Sto, d)**, with  $d \geq 1$ . It remains to precise  $d$ .

- *Step 2:*

*Case 2a:* If a **(Sto, d)** model is detected in Step 1, differentiate the current time series, and apply OPP test. Iterate this step until the null is rejected. Then  $d$  corresponds to the number of necessary differentiations.

*Case 2b:* If a **(Det, d)** model is detected in Step 1, differentiate the current time series, and apply KPSS test. Iterate this step until the null is not rejected. Then  $d$  corresponds to the number of necessary differentiations.

When applying this strategy, illustrated by a diagram in Figure D1, Appendix D, a model is suggested, leading either to equation (1) or (2), with parameters to be determined. In particular, process  $(B_t)_t$  is rarely a white noise, and should rather be modelled by an ARMA(p, q) process. The validity of the global model has to be confirmed with residuals diagnosis.

### 2.2.6 Periodic trend

To deal with seasonal data, we assume that the season  $r$  is known and will be equal to 12, 4 or 2 for monthly, quarterly or biannual data respectively. In this case we follow the Box et al.'s (1976) approach:

- at the first step, we remove the seasonal component
- in the second step, we apply our TDT strategy to the deseasonalised data.

In the first step one need to determine if the seasonal component is deterministic or stochastic, to consider the model (3) or (4). In Canova and Hansen (1995), the authors proposed the CH test where the null hypothesis is the presence of deterministic seasonal component whereas the alternative hypothesis is the presence of seasonal unit root. Hence we suggest to apply the CH test to the original data. If the null is not rejected, then we remove the seasonal component by using the harmonic regression to identify  $S_t$ , in (3) and the deseasonalised data will be  $Y_t = Z_t - S_t$ . If the null is rejected then the deseasonalised data will be  $Y_t = (1 - L^r)Z_t$ . We then apply our TDT strategy to  $Y_t$ .

## 3 Simulations

We ran 5,000 simulations using every DGP among **(SN)**, **(Det, 1)**, **(Det, 2)**, **(Sto, 1)** and **(Sto, 2)**, as defined from equations (6) to (10). We set  $n = 300$ ,  $a_0 = 5$ ,  $a_1 = 1$ ,  $a_2 = 1$ . Random generations of  $\mathcal{E}_t$  were taken from Gaussian centred variables with standard deviation  $\sigma_{\mathcal{E}}$ , a fixed value among  $\{0.5, 1, 3, 5, 10, 20, 30, 50, 100, 200, 300, 500\}$ , and the associated stationary noise is either a **(WN)**, or **(SN)** such as a MA(2) or an ARMA(1, 1) centred, causal and invertible process. Details on simulations are given in Supplementary S1.1.

### 3.1 Dickey-Fuller tests failure

Table 2 shows that Dickey-Fuller diagnosis is accurate for **(WN)** and **(Sto<sub>W</sub>, 1)** models. As expected, only the convenient model **(M<sub>3</sub>)**, provides a correct answer for **(Det<sub>W</sub>, 1)**. And in Figure E1 from Appendix E, we illustrate that diagnosis is unclear for **(Det<sub>W</sub>, 1)** realisations when using any test under models **(M<sub>1</sub>)** or **(M<sub>2</sub>)**. Indeed, results vary greatly according to parameter  $\sigma_{\mathcal{E}}$ , taking value in the set  $\{0.5, 1, 3, 5, 10, 20, 30, 50, 100, 200, 300, 500\}$ . Surprisingly, unit root is far to be correctly detected in **(Sto<sub>W</sub>, 2)** realisations. Finally, diagnosis is mainly incorrect for model **(Det<sub>W</sub>, 2)**, since unit roots are systematically detected, whereas we have  $\rho = 0$ . Thus, Dickey-Fuller-based tests fail in diagnosing unit root for quadratic trends models.

**Table 2** Null-hypothesis rejection rate (%) in Dickey-Fuller tests, when  $\sigma_{\mathcal{E}} = 10$

Test	DGP <sup>a</sup>				
	(WN)	(Det <sub>W</sub> , 1)	(Det <sub>W</sub> , 2)	(Sto <sub>W</sub> , 1)	(Sto <sub>W</sub> , 2)
Test $\rho$ under <b>(M<sub>1</sub>)</b>	100	0	0	5.06	0
Test $\rho$ under <b>(M<sub>2</sub>)</b>	100	0	0	4.96	12.88
Test $\rho$ under <b>(M<sub>3</sub>)</b>	100	100	0	5.04	32.4
Test $\Phi_1$ under <b>(M<sub>2</sub>)</b>	100	0	100	4.86	95.46
Test $\Phi_2$ under <b>(M<sub>3</sub>)</b>	100	100	100	4.92	99.08
Test $\Phi_3$ under <b>(M<sub>3</sub>)</b>	100	100	100	5.1	90.4

Note: <sup>a</sup>DGP.

Several strategies based on Dickey-Fuller tests have been developed (Dickey et al., 1986; Perron, 1988; Dolado et al., 1990). From Table 2, we can deduce that all the strategies permit to discriminate between first-order deterministic or stochastic trend, but they do not plan to integrate second-order trends. Thus, the most simple strategy proposed in Dickey et al. (1986) falsely classifies **(Det<sub>W</sub>, 2)** model as a random walk, whereas Perron (1988) predicts a linear-trend stationary process. The most advanced strategy given in Dolado et al. (1990) nearly identifies **(Det<sub>W</sub>, 2)** processes, by describing them as  $\Delta(Z_t) = \beta_0 + \beta_1 t + \mathcal{E}_t$ , instead of  $\Delta(Z_t) = \beta_0 + \beta_1 t + \Delta(\mathcal{E}_t)$ , where  $\Delta(Z_t) = Z_t - Z_{t-1}$  is the differentiated series. But diagnosis is mainly incorrect for **(Sto<sub>W</sub>, 2)** processes. Surprisingly, only the most simple strategy suggested in Dickey et al. (1986) detects unit root in half realisations, and predicts either a linear-trend stationary process or a stationary AR(1) process otherwise. And other strategies predict linear-trend stationary process. Consequently, it appears necessary to elaborate a new strategy to identify not only one-degree trends, but also higher-degree ones.

### 3.2 Accurate behaviour for OPP and KPSS tests

Table 3 presents the null-hypothesis rejection rate when  $B_t = \mathcal{E}_t$  where  $\sigma_{\mathcal{E}}$  varies in  $\{0.5, 1, 3, 5, 10, 20, 30, 50, 100, 200, 300, 500\}$ . It shows that KPSS and OPP tests perform as expected, not only for simulations driven from one-order trends, but also for quadratic trends. Moreover, results remain identical for any value of parameter  $\sigma_{\mathcal{E}}$ , varying from 0.5 to 500, such as shown in Figure F1, Appendix F. We also present results for simulations with underlying stationary noises  $(B_t)_t$ , such as a MA(2) and

an ARMA(1, 1). In this case, both Figures S1 and S2, in Supplementary S1.2, show the same behaviour for KPSS and OPP tests, whatever  $\sigma_{\mathcal{E}}$  value.

**Table 3** Null-hypothesis rejection rate (%) for KPSS and OPP stationarity tests, for DGP simulations when the underlying process is a white noise  $(\mathcal{E}_t)_t$

Test	DGP <sup>a</sup>				
	(WN)	(Det <sub>W</sub> , 1)	(Det <sub>W</sub> , 2)	(Sto <sub>W</sub> , 1)	(Sto <sub>W</sub> , 2)
OPP for $Z_t$	100	100	100	5.96	0
OPP for $\Delta(Z_t)$	100	100	100	100	5.86
KPSS for $Z_t$	4.8	98.6	100	98.9	100
KPSS for $\Delta(Z_t)$	0	0	100	4.8	98.9

Notes: <sup>a</sup>DGP.

We vary  $\sigma_{\mathcal{E}}$  on the set  $\{0.5, 1, 3, 5, 10, 20, 30, 50, 100, 200, 300, 500\}$ .

The final rejection rate is computed as the average of the rejection rates obtained for each  $\sigma_{\mathcal{E}}$ .

### 3.3 TDT strategy

We applied successively tests 1 to 4 to any simulation, using a risk  $\alpha = 5\%$ . In Table 4, every column returns the diagnosis percentage associated to the corresponding DGP. All the results with  $\sigma_{\mathcal{E}}$  value in the set  $\{0.5, 1, 3, 5, 10, 20, 30, 50, 100, 200, 300, 500\}$  are pooled, since the observed percentages show a very high stability with respect to  $\sigma_{\mathcal{E}}$ . As we have pointed out before, the percentage in bold refers to the theoretical diagnosis, i.e., the responses given in equations (11)–(15). We observe that, even if alternative responses can be observed, the modal responses given by the TDT strategy are the theoretical ones.

Let us first consider simulations generated from a (WN) model, only two sets of responses were obtained to TDT: **Alt/Alt/Null/Null** for 95.25% of (WN) simulations (percentage written in bold), and **Alt/Alt/Alt/Null** for the other 4.75% simulations. This result is convenient with the expected behaviour of KPSS test, as described in Subsection 2.2.5.2, equation (16), where the expected percentage is 95% (resp. 5%) for **-/-/Null/Null** (resp. **-/-/Alt/Null**) response. The new element that appears thanks to the simulations is that OPP test is consistent for a stationary noise, i.e., the power converges to one as  $n \rightarrow +\infty$ . As far as (Det<sub>W</sub>, 1) or (Det<sub>W</sub>, 2) processes are concerned, almost only the theoretical response was observed (99.99% and 100% resp.), which is in line with the expected results given in equations (17) and (18). Once again, the novelty lies in the high power of the OPP test for series generated by a deterministic trend, as defined in equation (1). Finally, for (Sto<sub>W</sub>, 1) or (Sto<sub>W</sub>, 2) simulations, more responses are observed, but they remain consistent with the expected results presented in Subsection 2.2.5.2. On the one hand, (Sto<sub>W</sub>, 1) provides observed response rates very close to the expected ones, given in equation (19). Indeed the observed percentage 89.38% (resp. 4.28%, 5.91% and 0.44%) for **Null/-/-/Null** (resp. **Null/-/-/Alt**, **Alt/-/-/Null** and **Alt/-/-/Alt**) response is very close to the expected one 90.25% (resp. 4.75%, 4.75% and 0.25%). This time, to compute the observed percentages, it is necessary to sum over the convenient associated responses. For instance, the 89.38% rate for the observed **Null/-/-/Null** response is the sum of percentages 86.162% and 3.216% observed for **Null/Alt/Null/Null** and **Null/Null/Alt/Null** responses respectively.

On the other hand (**Sto<sub>W</sub>, 2**) simulations show that KPSS test has a high power on these series, detecting almost always the non-stationarity induced by a double unit root, and moreover that even if OPP test was not developed to deal with multiple unit root, it detects, in a too conservative way, the presence of this double unit root.

But the interest of Table 4 lies is in its reverse reading. Let us associate a DGP to a TDT diagnosis. As an example, since none simulation under (**WN**), (**Det<sub>W</sub>, 1**), (**Det<sub>W</sub>, 2**) or (**Sto<sub>W</sub>, 1**) DGP led to responses **Null/Null/Alt/Alt**, then if one obtains such a response on its time series, this means that a (**Sto<sub>W</sub>, 2**) model is suitable. Alternately **Alt/Alt/Alt/Null** diagnosis could lead either to a (**WN**), a (**Det<sub>W</sub>, 1**) or a (**Sto<sub>W</sub>, 1**) model. But referring to occurrence percentages,  $99.993/(4.753 + 99.993 + 5.822) = 90.436\%$  of simulations with TDT **Alt/Alt/Alt/Null** are produced by a (**Det<sub>W</sub>, 1**) DGP, only 4.299% by a (**WN**) and 5.266% by a (**Sto<sub>W</sub>, 1**). Then a (**Det<sub>W</sub>, 1**) model appears as the best candidate, but (**WN**) and (**Sto<sub>W</sub>, 1**) models cannot be totally excluded. In this precise case, even if a linear trend is present, the relevance of either a (**WN**) or a (**Det<sub>W</sub>, 1**) model depends on the intensity of the trend in relation to the variance of the associated noise. The convenient choice can be ruled out by previously computing autocorrelation functions. If several models remain acceptable, then we suggest to construct and compare them.

**Table 4** Percentage of TDTs associated to every DGP when  $\sigma_{\varepsilon}$  takes values in  $\{0.5, 1, 3, 5, 10, 20, 30, 50, 100, 200, 300\}$

TDT <sup>b</sup>	DGP <sup>a</sup>				
	(WN)	(Det <sub>W</sub> , 1)	(Det <sub>W</sub> , 2)	(Sto <sub>W</sub> , 1)	(Sto <sub>W</sub> , 2)
Alt/Alt/Alt/Alt	0	0	100 <sup>c</sup>	0.438	0
Alt/Alt/Null/Alt	0	0	0	0	0
Alt/Null/Alt/Alt	0	0	0	0	0
Alt/Null/Null/Alt	0	0	0	0	0
Null/Alt/Alt/Alt	0	0	0	4.273	6.213
Null/Alt/Null/Alt	0	0	0	0.005	0
Null/Null/Alt/Alt	0	0	0	0	<i>90.427</i>
Null/Null/Null/Alt	0	0	0	0	0.007
Alt/Alt/Alt/Null	4.753	<i>99.993</i>	0	5.822	0
Alt/Alt/Null/Null	<i>95.247</i>	0.007	0	0.084	0
Alt/Null/Alt/Null	0	0	0	0	0
Alt/Null/Null/Null	0	0	0	0	0
Null/Alt/Alt/Null	0	0	0	<i>86.162</i>	0.089
Null/Alt/Null/Null	0	0	0	3.216	0
Null/Null/Alt/Null	0	0	0	0	3.220
Null/Null/Null/Null	0	0	0	0	0.015
Total percentage	100	100	100	100	100

Notes: <sup>a</sup>DGP.

<sup>b</sup>TDTs.

<sup>c</sup>Italic font highlights the expected TDT diagnosis associated to every DGP.



Let us remark that classification stability may decrease when  $\sigma_{\mathcal{E}}$  keeps growing. Indeed when noise intensity becomes too high compared to the linear coefficient  $a_1$ , the trend becomes imperceptible, and KPSS test may progressively fail to reject the null for several **(Det<sub>W</sub>, 1)** simulations. Then the accurate response **Alt/Alt/Alt/Null** may successively be replaced by the wrong response **Alt/Alt/Null/Null**, that is accurate for **(WN)**. And the confusion between **(Det<sub>W</sub>, 1)** and **(WN)** naturally increases with  $\sigma_{\mathcal{E}}$ . But note that in this case, the true model **(Det<sub>W</sub>, 1)** might no longer be the most suitable for the series, and a simple **(WN)** model should be preferred.

In order to evaluate the effect of autocorrelation on TDT strategy, we also run simulations with autocorrelated noises. In other words, we replaced  $\mathcal{E}_t$  either by a MA(2) process or by an ARMA(1, 1) process, denoted as  $B_t$ , see Supplementary S1.3. Percentage of TDT diagnosis associated to every DGP, driven by **(SN)**, are given in Supplementary, in Tables S1 and S2. Percentage diagnosis remains similar regardless of the underlying model for  $B_t$ , and are still highly stable with respect to  $\sigma_{\mathcal{E}}$  value. From Table 4, S1 and S2, we suggest to associate each model with some sets of responses to TDT.

Finally, from Table 4 we observe that the five responses **Alt/Alt/Null/Alt**, **Alt/Null/Alt/Alt**, **Alt/Null/Null/Alt**, **Alt/Null/Alt/Null** and **Alt/Null/Null/Null** never appear as output of the TDT strategy, therefore we propose the following associations:

- **Alt/Alt/Null/Null** is associated with **(SN)**,
- **Alt/Alt/Alt/Null** is associated with **(Det, 1)**,
- **Alt/Alt/Alt/Alt** is associated with **(Det, 2)**,
- **Null/Alt/Alt/Null** are associated with **(Sto, 1)**,
- Null/Alt/Null/Null**
- Null/Alt/Null/Alt** (20)
- **Null/Null/Alt/Alt** are associated with **(Sto, 2)**,
- Null/Null/Alt/Null**
- Null/Null/Null/Alt**
- Null/Null/Null/Null**
- **Null/Alt/Alt/Alt** is associated either with **(Sto, 1)** or **(Sto, 2)**.

## 4 Application on data

### 4.1 Money stock in the USA


We consider money stock evolution in USA, that is given in billions of dollars and annual averaged from 1889 to 1988 [see Figure 3(a)]. Autocorrelation functions, given in Figure 3(b), confirm that data are driven by a trend.

Table 5 shows that our strategy TDT applied to money stock data suggests a **(Sto, 2)** model. Dickey-Fuller tests rather suggest a **(Sto, 1)** model. Indeed, most Dickey-Fuller tests do not reject the null for the initial series, but they are unable to detect a unit root in the differentiated series. We construct both a **(Sto, 1)** and a **(Sto, 2)** model, as in equation (2), by modelling the stationary process  $(B_t)_t$  by the first clearly valid model among all ARMA( $p, q$ ) with  $p, q \leq 2$ , sorted by minimising Schwarz's Bayesian criterion (Schwarz, 1978):

- $(B_t)_t$  is MA(2) for **(Sto, 2)** model, suggested by our strategy TDT
- $(B_t)_t$  is ARMA(1, 1) for **(Sto, 1)** model, suggested by Dickey-Fuller tests.

In Table 6, we compare both models relevance in terms of prediction criteria computed between the observed series and predictions for the last 10 values (almost 10%), such as root mean square error (RMSE) and mean absolute percentage error (MAPE). Table 6 shows that the **(Sto, 2)** model provided by our strategy is the most suitable. In Supplementary S2, we additionally explore the other US macroeconomic indexes contained in Nelson-Plosser data.

**Figure 3** (a) Money stock data evolution (b) Autocorrelation functions (see online version for colours)



**Table 5** p-values provided by several tests on the initial and the differentiated money stock series


Series	Test				
	OPP	KPSS	$\rho$ under $(M_1)$	$\rho$ under $(M_2)$	$\rho$ under $(M_3)$
$Z_t$	0.2	0.01	0.99	0.99	0.94
$\Delta(Z_t)$	0.09	0.1	0.01	0.01	0.01

**Table 6** Models comparison for money stock series

Model	Criterion	
	RMSE	MAPE
TDT <sup>a</sup> : <b>(Sto, 2)</b> with MA(2)	0.038	0.362
DF <sup>b</sup> : <b>(Sto, 1)</b> with ARMA(1, 1)	0.317	3.549


Notes: <sup>a</sup>TDTs.

<sup>b</sup>Dickey-Fuller tests.

**Figure 4** Predictions for money stock series with TDT and DF models (see online version for colours)

#### 4.2 Evolution of atmospheric $\text{CO}_2$ concentration

Since 1959, atmospheric  $\text{CO}_2$  concentration (ppm) has been measured monthly, at Mauna Loa Observatory, Hawaii, see Keeling et al. (2001). Figure 5 reveals that the global average concentration of atmospheric carbon dioxide has a clear increasing trend, and also a seasonal monthly component.


**Figure 5**  $\text{CO}_2$  atmospheric concentration evolution

As described in Subsection 2.2.6, we start by removing the seasonal component. Since the data are given monthly, the underlying period is  $r = 12$ . CH-test indicates that the seasonality is deterministic (CH-test, p-value = 1). Thus we deseasonalise, by regressing the  $\text{CO}_2$  series on the seasonal dummy variables and by retaining the residuals from this regression. Autocorrelation functions plotted in Figure 6(b) show that the remaining series is driven by a trend. From Table 7, we see that Dickey-Fuller tests applied to the deseasonalised series clearly suggests a **(Sto, 1)** model, whereas our TDT strategy produces responses **Alt/Alt/Alt/Alt** to tests 1 to 4, suggesting rather a **(Det, 2)** model. We construct both a **(Sto, 1)** and a **(Det, 2)** model, as in equations (1) and (2), by

modelling the stationary process  $(B_t)_t$  by the first valid model among all SARMA( $p, q$ ) ( $P, Q$ ) (Dolado et al., 1990) with  $p, q, P, Q \leq 2$ , sorted by minimising Schwarz's Bayesian criterion (Schwartz, 1978):

- $(B_t)_t$  is SARMA(1, 2)(1, 1) (Dolado et al., 1990) for **(Det, 2)** model, suggested by our strategy TDT.
- $(B_t)_t$  is SARMA(1, 0)(1, 1) (Dolado et al., 1990) for **(Sto, 1)** model, suggested by Dickey-Fuller tests.

**Figure 6** (a) CO<sub>2</sub> deseasonalised series evolution (b) Associated autocorrelation functions (see online version for colours)



In Table 8, we compare both models relevance in terms of information criteria such as AIC, BIC, AICc (Akaike, 1973; Schwartz, 1978; Hurvich and Tsai, 1991) and prediction criterion computed between the observed series and predictions for the last six years (almost 10%), such as RMSE and MAPE. Table 8 shows that the **(Det, 2)** model provided by our strategy is the most suitable.

**Table 7** p-values provided by several tests on CO<sub>2</sub> deseasonalised series

Series	Test				
	OPP	KPSS	$\rho$ under ( $M_1$ )	$\rho$ under ( $M_2$ )	$\rho$ under ( $M_3$ )
$Z_t$	0.01	0.01	0.9791	0.99	0.5465
$\Delta(Z_t)$	0.01	0.01	0.01	0.01	0.01

**Table 8** Models comparison for CO<sub>2</sub> deseasonalised series


Model	Criterion	
	RMSE	MAPE
TDT <sup>a</sup> : <b>(Det, 2)</b> with SARMA(1, 2)(1, 1) (Dolado et al., 1990)	0.78	1.291
DF <sup>b</sup> : <b>(Sto, 1)</b> with SARMA(1, 0)(1, 1) (Dolado et al., 1990)	1.757	2.816

Notes: <sup>a</sup>TDTs.

<sup>b</sup>Dickey-Fuller tests.

Figure 7 shows that forecasts for CO<sub>2</sub> atmospheric concentration maintain the same trajectory, with great accuracy. Indeed, prediction intervals are so thin that they are hardly visible.

**Figure 7** Predictions for CO<sub>2</sub> atmospheric concentration with TDT model, where prediction intervals coloured in steel blue (resp. light grey) represent 80% (resp. 95%) confidence level (see online version for colours)



## 5 Conclusions

We present a strategy to detect and identify the nature of the trend component in time series. We recall that, as a first analysis, visualising time series plot is indispensable, since it may already suggest the presence of a trend. Then trend can be confirmed by analysing autocorrelation functions. Next, whence a trend is detected, it remains to identify its nature. Indeed deterministic or stochastic trends do not produce the same forecasts.

We have shown that the Dickey-Fuller test is not suitable. For instance it detects the presence of spurious unit root for series generated under model **(Det<sub>W</sub>, 2)**. In this paper, we rather propose a strategy based on OPP and KPSS tests, called TDT, in order to select between either a **(Det, d)** (i.e., a deterministic polynomial trend) or a **(Sto, d)** (i.e., a stochastic trend). We have given theoretical justifications on the expected results when applying our TDT strategy to data (Subsections 2.2.5.1 and 2.2.5.2). In particular, we had to study the asymptotic behaviour of the KPSS test statistic when the required assumptions are not verified [either a non-invertible moving average or the two alternatives **(Det<sub>W</sub>, 1)** and **(Det<sub>W</sub>, 2)**]. Next our TDT strategy was assessed on simulations. We have performed simulations by considering some particular cases of these models namely **(WN)** (i.e. a white noise), **(Det<sub>W</sub>, 1)** (i.e., a deterministic linear trend), **(Det<sub>W</sub>, 2)** (i.e., a deterministic quadratic trend), **(Sto<sub>W</sub>, 1)** (i.e., a stochastic trend with a single unit root) and **(Sto<sub>W</sub>, 2)** (i.e., a stochastic trend with a double unit root). It appears that the results observed on the simulations (Table 4) correspond to the expected results. Moreover, the simulations allow us to discuss the power of the OPP test when it is applied under the alternative hypothesis, and confirm that KPSS test has a size

distortion if the underlying model is a non-invertible moving average, since it never rejects the null. We have also pointed out that KPSS test is consistent under the two alternatives (**Det<sub>W</sub>, 1**) and (**Det<sub>W</sub>, 2**), since it always rejects the null in this case.

Let us remark that both procedures, TDT strategy and Dickey-Fuller tests, may result in the same model suggestion. But diagnosis can be different, especially for time series with a quadratic trend, that reduces Dickey-Fuller tests reliability. Hence when both procedures suggest different models, both model candidates should be computed, validated and compared. Applied on two real datasets in Section 4, our automatic TDT strategy provides a better model than the one provided using the classical Dickey-Fuller tests. Thus we illustrated the superiority of our strategy on the Dickey-Fuller tests in selecting the best models which minimise the error of prediction criteria, such as the RMSE and the MAPE. Therefore we recommend to use our TDT strategy in modelling non-stationary time series as a procedure to choose a candidate model with an accurate nature of the trend.

As a conclusion, applying our strategy on data allows to identify a candidate model, which must be validated and eventually compared with other potential models.

Appendices/Supplementary materials are available on request by emailing the corresponding author or can be obtained under <http://www.i2m.univ-amu.fr/perso/manuela.royer-carezni/AnnexesR.TrendTS/TrendTS.html>.

## References

- Akaike, H. (1973) 'Information theory as an extension of the maximum likelihood principle', in Petrov, B.N. and Csaki, F. (Eds.): *Second International Symposium on Information Theory*, Vol. 1557, Springer, Akademiai Kiado, Budapest, Hungary, pp.30–32.
- Boutahar, M. (2009) 'Comparison of non-parametric and semi-parametric tests in detecting long memory', *Journal of Applied Statistics*, Vol. 36, No. 9, pp.945–972.
- Box, G.E.P., Jenkins, G.M. and Reinsel, G.C. (1976) *Time Series Analysis: Forecasting and Control*, 3rd ed., Holden-Day, San Francisco.
- Brockwell, P.J. and Davis, R.A. (1991) *Time Series: Theory and Methods*, 2nd ed., Springer-Verlag, New York.
- Canova, F. and Hansen, B.E. (1995) 'Are seasonal patterns constant over time? A test for seasonal stability', *Journal of Business and Economic Statistics*, Vol. 13, No. 3, pp.237–252.
- Chan, N.H. and Wei, C.Z. (1988) 'Limiting distributions of least squares estimates of unstable autoregressive processes', *Ann. Statist.*, Vol. 16, No. 1, pp.367–401.
- Chan, K.H., Hayya, J.C. and Ord, J.K. (1977) 'A note on trend removal methods: the case of polynomial regression versus variate differencing', *Econometrica*, Vol. 45, No. 3, pp.737–744.
- Conway, J.H. and Guy, R.K. (1996) *The Book of Numbers*, Springer-Verlag, New York.
- Davydov, Y.A. (1970) 'The invariance principle of stationary processes', *Theory of Probability and its Application*, Vol. 15, No. 3, pp.487–498.
- Dickey, D.A. and Fuller, W.A. (1979) 'Distribution of the estimators for autoregressive time series with a unit root', *J. Am. Stat. Assoc.*, Vol. 74, No. 366, pp.427–431.
- Dickey, D.A., Bell, W.R. and Miller, R.B. (1986) 'Unit roots in time series models: tests and implications', *J. Am. Stat. Assoc.*, Vol. 40, No. 1, pp.12–26.
- Dolado, J.J., Jenkinson, T. and Sosvilla-Rivero, S. (1990) 'Cointegration and unit roots', *J. Econ. Surv.*, Vol. 4, No. 3, pp.249–273.

- Ertur, C. (1998) *Méthodologies de test de la racine unitaire*, Rapport de recherche, Laboratoire d'Analyses et de techniques économiques (LATEC), Université de Bourgogne, hal-01527262.
- Hardy, G.H. and Wright, E.M. (1975) *An Introduction to the Theory of Numbers*, 4th with corrections ed., Springer-Verlag, Clarendon Press, Oxford.
- Hurvich, C.M. and Tsai, C-L. (1991) 'Bias of the corrected AIC criterion for underfitted regression and time series models', *Biometrika*, Vol. 78, No. 3, pp.499–509.
- Keeling, C.D., Piper, S.C., Bacastow, R.B., Wahlen, M., Whorf, T.P., Heimann, M. and Meijer, H.A. (2001) 'Exchanges of atmospheric CO<sub>2</sub> and <sup>13</sup>CO<sub>2</sub> with the terrestrial biosphere and oceans from 1978 to 2000. I. Global aspects', *SIO Reference Series*, Vol. 6, No. 1, pp.1–28.
- Kwiatkowski, D., Phillips, P.C.B., Schmidt, P. and Shin, Y. (1992) 'Testing the null hypothesis of stationarity against the alternative of a unit root', *Journal of Econometrics*, Vol. 54, Nos. 1–3, pp.159–178.
- Nelson, C.R. and Kang, H. (1981) 'Spurious periodicity in inappropriately detrended time series', *Econometrica*, Vol. 49, No. 3, pp.741–751.
- Nelson, C.R. and Plosser, C. (1982) 'Trends and random walks in macroeconomics time series: some evidences and applications', *Journal of Monetary Economics*, Vol. 10, No. 2, pp.139–162.
- Ouliaris, S., Park, J.Y. and Phillips, P.C.B. (1989) 'Testing for a unit root in the presence of a maintained trend', *Advances in Econometrics and Modeling*, Kluwer Academic Publishers, Needham, MA.
- Perron, P. (1988) 'Trends and random walk in macroeconomics time series: further evidence from a new approach', *JEDC*, Vol. 12, Nos. 2–3, pp.297–332.
- Phillips, P.C.B. and Perron, P. (1988) 'Testing for a unit root in time series regression', *Biometrika*, Vol. 75, No. 2, pp.335–346.
- Prokhorov, Y.V. (1956) 'Convergence of random processes and limit theorems in probability theory', *Theory Probab. Appl.*, Vol. 1, No. 2, pp.157–214.
- Schwartz, G.E. (1978) 'Estimating the dimension of a model', *Ann. Stat.*, Vol. 6, No. 2, pp.461–464.